


Datasheet for IR44-131 anti-Huntingtin antibody		Package size: 25, 100 μl Store at: -20°C MW (kDa): 350
Overview		
Product Name	Huntingtin antibody	
Product Number	IR44-131	
Gene Description	huntingtin	
Clonality	Polyclonal	
Host	Rabbit	
Species Reactivity	Human , mouse, rat	
Recommended Applications Dilutions	Western Blot 1:1000 - 1:2000 Immunofluorescence 1:200 – 1:500	
Storage Buffer	100mM Tris Glycine, 20% Glycerol (pH7). 0.025% ProClin 300 was added as a preservative	
Concentration	1.25 mg/ml	
Purity	Affinity column purified	
Storage	Store at +4°C for short term storage. Long time storage is recommended at -20°C	
Notes	Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user.	
Image		
		IR44-131 anti-Huntingtin antibody WB image
		<p>All lanes : Anti-Huntingtin antibody at 1/1000 dilution</p> <p>Lysates/proteins at 60 μg per lane This blot was produced using a 5% SDS-PAGE. The gel was run at 140V for 50 minutes before being transferred onto a Nitrocellulose membrane at 18V for 60 minutes. The membrane was then blocked for an hour before being incubated with IR44-131 overnight at 4°C.</p>






IReal Biotechnology Co., Ltd.

7F., No. 9, Sec. 3, Zhonghua Rd., North Dist., Hsinchu City 300, Taiwan (R.O.C.)

Website : www.irealbio.com

TEL : 03-526-0005

E-mail : irealbio@irealbio.com

Customer feedback Image	
	<p>IR44-131 anti-Huntingtin antibody WB image</p> <p>Sample : Neuron 2A Huntingtin (IR44-131) : 1-1000 Data from : Taipei Medical University</p>
Image	
<p style="text-align: center;">Huntingtin</p> <div style="display: flex; flex-direction: column; align-items: center;"> <div style="display: flex; justify-content: space-around; width: 100%;"> <div style="writing-mode: vertical-rl; transform: rotate(180deg);">DBTRG-05MG</div>   </div> <div style="display: flex; justify-content: space-around; width: 100%;"> <div style="writing-mode: vertical-rl; transform: rotate(180deg);">HUVEC</div>   </div> </div>	<p>IR44-131 anti-Huntingtin antibody ICC/IF image</p> <p>Immunofluorescence: cells were fixed with 4% paraformaldehyde for 10 min at RT, permeabilized with 0.1% NP-40 for 10 min at RT then blocked with 5% BSA for 30 min at room temperature. Cells were stained with IR44-131 anti-Huntingtin antibody (red) at 1:200 and 4°C. DAPI (blue) was used as the nuclear counter stain.</p>

IReal Biotechnology Co., Ltd.

7F., No. 9, Sec. 3, Zhonghua Rd., North Dist., Hsinchu City 300, Taiwan (R.O.C.)

Website : www.irealbio.com

TEL : 03-526-0005

E-mail : irealbio@irealbio.com

