

Catalog Number: IR136-543


Anti-NRF2 antibody

Package size: 25, 100 μ l
Store at: -20°C
MW (kDa): 125-160


Overview

Product Name	NRF2 antibody
Product Number	IR136-543
Gene Description	nuclear factor, erythroid 2-like 2
Clonality	Polyclonal
Host	Rabbit
Species Reactivity	Human, Mouse, Rat
Recommended Applications Dilutions	Western Blot 1:500 – 1:1000 Immunofluorescence 1:100 – 1:200 Immunohistochemistry (Paraffin) 1:100 – 1:200
Storage Buffer	100mM Tris Glycine, 20% Glycerol (pH7). 0.025% ProClin 300 was added as a preservative
Concentration	1.41 mg/ml
Purity	Affinity column purified
Storage	Store at +4°C for short term storage. Long time storage is recommended at -20°C
Notes	Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user.

IR136-543 anti-NRF2 antibody WB image



IR136-543 anti-NRF2 antibody IHC image



Anti-NRF2 antibody at 1/800 dilution.
Lysates at 30 μ g per lane. This blot was produced using a 5% SDS-PAGE. NC membrane was then blocked for an hour before being incubated with IR136-543 overnight at 4°C.

Immunohistochemical analysis of paraffin embedded Human cancer tissue labeling NRF2 with IR136-543 at 1/150.

Image	
<div> <div>NRF2</div> <div> <div>HT-29</div> <div>MDA-MB-231</div> </div> </div>	<div>IR136-543 anti-NRF2 antibody ICC/IF image</div> <div> <p>Immunofluorescence: cells were fixed with 4% paraformaldehyde for 10 min at RT, permeabilized with 0.1% NP-40 for 10 min at RT then blocked with 5% BSA for 30 min at room temperature. Cells were stained with IR136-543 anti-NRF2 antibody (red) at 1:200 and 4°C. DAPI (blue) was used as the nuclear counter stain.</p> </div>
Image	
	<div>IR136-543 anti-NRF2 antibody IHC image</div> <div> <p>Immunohistochemical analysis of paraffin embedded mouse tissue labeling NRF2 with IR136-543 at 1/100 RT 1hr. Epitope Retrieval methods: Citrate Buffer, pH 6.0</p> <p>Data from: Li-Tzung Biotechnology. Co</p> </div>

IReal Biotechnology Co., Ltd.

