

Catalog Number: IR48-138


Anti-Fibronectin (C-term) antibody

Package size: 25, 100 μ l
Store at: -20°C
MW (kDa): 130-300


Overview

Product Name	Fibronectin (C-term) antibody
Product Number	IR48-138
Gene Description	fibronectin 1
Clonality	Polyclonal
Host	Rabbit
Species Reactivity	Human, mouse, rat
Recommended Applications Dilutions	Western Blot 1:500 Immunofluorescence 1:200 – 1:500 Immunohistochemistry (Paraffin) 1:100 – 1:200
Storage Buffer	100mM Tris Glycine, 20% Glycerol (pH7). 0.025% ProClin 300.
Concentration	1.5 mg/ml
Purity	Affinity column purified
Storage	Store at +4°C for short term storage. Long time storage is recommended at -20°C
Notes	Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user.

IR48-138 anti-Fibronectin antibody WB image



IR48-138 anti-Fibronectin antibody IHC image




Anti-Fibronectin antibody at 1/500 dilution. Lysates at 60 μ g per lane. This blot was produced using a 5% SDS-PAGE. Nitrocellulose membrane was then blocked with 3%BSA for an hour before being incubated with IR48-138 overnight at 4°C.

Immunohistochemical analysis of paraffin embedded human hepatocellular carcinoma (HCC) labeling Fibronectin with IR48-138 at 1/100.


IR48-138 anti-Fibronectin antibody IHC image

Human Breast Carcinoma



Immunohistochemical analysis of paraffin embedded human breast carcinoma labeling Fibronectin with IR48-138 at 1/100.


IR48-138 anti-Fibronectin antibody IHC image



Immunohistochemical analysis of paraffin embedded Human cancer tissue labeling Fibronectin with IR48-138 at 1/100.


IR48-138 anti-Fibronectin antibody ICC/IF image

PANC-1 cells



Immunofluorescence: cells were fixed with 4% paraformaldehyde for 10 min at RT, permeabilized with 0.1% NP-40 for 10 min at RT then blocked with 5% BSA for 30 min at room temperature. Cells were stained with IR48-138 anti-Fibronectin antibody (red) at 1:200 and 4°C. DAPI (blue) was used as the nuclear counter stain.

IR48-138 anti-Fibronectin antibody WB image



Anti-Fibronectin antibody at 1/1000 dilution. Lysates at 50 µg per lane. This blot was produced using a 5% SDS-PAGE. Nitrocellulose membrane was incubated with IR48-138 overnight at 4°C.
*IR292-962 Collagen I / COL1A1 antibody.

IReal Biotechnology Co., Ltd.