

WB

IF

IHC

Catalog Number: IR219-807


## Anti c-MET antibody

Package size: 25, 100  $\mu$ l  
Store at: -20°C  
MW (kDa): 130-170,  
p40 MET


### Overview

<b>Product Name</b>	c-MET antibody
<b>Product Number</b>	IR219-807
<b>Gene Description</b>	MET proto-oncogene, receptor tyrosine kinase
<b>Clonality</b>	Polyclonal
<b>Host</b>	Rabbit
<b>Species Reactivity</b>	Human, Mouse, Rat
<b>Recommended Applications Dilutions</b>	Western Blot 1:500 – 1:1000 Immunofluorescence 1:100 – 1:200 Immunohistochemistry (Paraffin) 1:100
<b>Storage Buffer</b>	100mM Tris Glycine, 20% Glycerol (pH7). 0.025% ProClin 300 was added as a preservative
<b>Concentration</b>	1.53 mg/ml
<b>Purity</b>	Affinity column purified
<b>Storage</b>	Store at +4°C for short term storage. Long time storage is recommended at -20°C
<b>Notes</b>	Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user.

### IR219-807 anti c-MET antibody WB image



### IR219-807 anti c-MET antibody IHC image




#### Anti c-MET antibody at 1/700 dilution.


Lysates at 50  $\mu$ g per lane. This blot was produced using a 5% SDS-PAGE. NC membrane was then blocked for an hour before being incubated with IR219-807 overnight at 4°C.

Immunohistochemical analysis of paraffin embedded Human Breast cancer tissue labeling c-MET with IR219-807 at 1/100.

**IR219-807 anti c-MET antibody WB image**



**IR219-807 anti c-MET antibody IHC image**




**Anti c-MET antibody at 1/1000 dilution.**


Lysates at 50 µg per lane. This blot was produced using a 5% SDS-PAGE. NC membrane was then blocked for an hour before being incubated with IR219-807 overnight at 4°C.

Immunohistochemical analysis of paraffin embedded Human cancer tissue labeling c-MET with IR219-807 at 1/100.

**IR219-807 anti c-MET antibody WB image**



**IR219-807 anti c-MET antibody IHC image**



**Anti c-MET antibody at 1/800 dilution.**

Lysates at 50 µg per lane. This blot was produced using a 5% SDS-PAGE. NC membrane was then blocked for an hour before being incubated with IR219-807 overnight at 4°C.

Immunohistochemical analysis of paraffin embedded Human Breast cancer tissue labeling c-MET with IR219-807 at 1/100.

IReal Biotechnology Co., Ltd.