

Catalog Number: IR301-983


Anti-CD63 antibody

Package size: 25, 100 μ l
Store at: -20°C
MW (kDa): 25-60


Overview

Product Name	CD63 antibody
Product Number	IR301-983
Gene Description	CD63 molecule
Clonality	Polyclonal
Host	Rabbit
Species Reactivity	Human, Mouse, Rat
Recommended Applications Dilutions	Western Blot 1:500 – 1:1000 Flow Cytometry 1:100 Immunohistochemistry (Paraffin) 1:200 – 1:400
Storage Buffer	100mM Tris Glycine, 20% Glycerol (pH7). 0.025% ProClin 300 was added as a preservative
Concentration	1.51 mg/ml
Purity	Affinity column purified
Storage	Store at +4°C for short term storage. Long time storage is recommended at -20°C
Notes	Gently mix before use. Optimal concentrations and conditions for each application should be determined by the user.

IR301-983 anti-CD63 antibody WB image




IR301-983 anti-CD63 antibody IHC image




at 1/1000 dilution. Lysates at 10 μ g per lane. This blot was produced using a 12% SDS-PAGE. Nitrocellulose membrane was then blocked for an hour before being incubated with IR301-983 overnight at 4°C.

Immunohistochemical analysis of paraffin embedded Human cancer tissue labeling CD63 with IR301-983 at 1/100.

IR301-983 anti-CD63 antibody WB image





IR301-983 anti-CD63 antibody FACS image



CD63 antibody at 1/1000 dilution. Lysates at 30 µg per lane. This blot was produced using a 12% SDS-PAGE. Nitrocellulose membrane was then blocked for an hour before being incubated with IR301-983 overnight at 4°C.

Product: CD63 antibody
 Catalog: IR301-983
 Cell: Human peripheral blood lymphocytes
 CD63 antibody: 1-100
Data from: National Chiao-Tung University

**Customer Feedback
 IR301-983 anti-CD63 antibody WB image**



CD63 antibody at 1/1000 dilution. Lysates at 50 µg per lane. This blot was produced using a 12% SDS-PAGE. Nitrocellulose membrane was then blocked for an hour before being incubated with IR301-983 overnight at 4°C.

CD63 antibody at 1/1000 dilution. Lysates at 50 µg per lane. This blot was produced using a 12% SDS-PAGE. Nitrocellulose membrane was then blocked for an hour before being incubated with IR301-983 overnight at 4°C.

IReal Biotechnology Co., Ltd.

